

Precautions & Prevention of Spread

1. Good hand washing, especially after handling tissues to wipe eyes or blow nose.



2. Proper disposal of tissues.

3. Separate towel and washcloth for the infected person.

4. Disinfect toys and other items handled by infected person. Wash in dishwashing detergent and rinse in a dilute bleach solution (one part bleach to ten parts water).



5. Other environmental surfaces in the child's play area, such as tables, may be cleaned with standard household cleaning materials or dilute bleach solution.



6. Towels and washcloths should be washed in hot water and dried in a hot dryer.

7. Avoid touching or rubbing the eyes.



All Client Information is kept confidential.



Administration	758-3820
HIV Testing & Counseling	758-3880
Communicable Disease	758-3885
Community Resource Nurse	758-3675
Dental	758-3835
Environmental Health	758-3830
Family Planning	758-3870
Hearing Impaired Line	758-3725
Hearing & Vision	758-3890
Immunization	758-3840
Laboratory	758-3825
Lead Testing	758-3768
Nursing Services	758-3850
Saginaw Health Plan	758-3855
Sexually Transmitted Diseases	758-3880
Substance Abuse Assessment	758-3781
After Hours	888-466-3141
WIC	758-3733

PINKEYE



Saginaw County
Department of Public
Health

1600 N. Michigan Ave.
Saginaw, MI 48602
(989) 758-3800

“Protecting and promoting the public’s health
and well-being”

www.saginawpublichealth.org



Pinkeye (Conjunctivitis)

An infection of the moist tissue that covers the front of the eyeball and lines the eyelids. It can be caused by a bacteria or virus.



Pinkeye can infect anyone, but it is most common in preschool and elementary school age children.

Mode of Spread

- ◆ Contact with discharges from eyes, nose or throat of infected people.
- ◆ Fingers, clothes, toys and other items contaminated with the discharges, including shared eye make-up applicators and multi-dose eye medications.



Incubation Period

Within 24 - 72 hours of contact with the discharges of the infected person.

Symptoms



1. Tearing of the infected eye.
2. Irritation and redness of the moist tissue that covers the eyeball and lines the eyelids. (This symptom is what gives Pinkeye its name).
3. Swelling of the affected eyelid.
4. Discharge from the affected eye that starts out clear yellow and quickly progresses to thick yellow pus and then green.

PINKEYE THAT STARTS OUT IN ONE EYE CAN QUICKLY SPREAD TO THE OTHER EYE!



Treatment



- Eye ointment or drops prescribed by the doctor are applied to the eye as directed.
- Be sure to wash your hands before inserting medication.



Exclusion from School

1. Exclude during the acute phase when puss is draining from the eyes.
2. Children with bacterial conjunctivitis may return to school after 24 hours of appropriate treatment.
3. Children with viral conjunctivitis should be excluded until all symptoms are resolved.

