

**Tips to Enhance Treatment
and Prevent Spread**

All client information is kept confidential.



1. Wash the sore thoroughly with soap and water to remove scabs or crusts.
2. Apply treatment as directed.
3. Wash your hands after applying Antifungal cream.
4. Towels, washcloths, bedding, and clothing that have been in contact with the infected person should be washed in hot water and dried in a hot dryer.
5. Use a separate towel and washcloth for the infected person.
6. Combs and brushes should be thoroughly cleaned by soaking in boiling hot water for ten minutes or by washing with any household disinfectant/detergent and hot water.
7. Do not share combs, brushes, hats, or other headgear.
8. Continue treatment for two weeks after it appears to be healed.
9. Ringworm infection is very persistent BUT DO NOT SKIP ANY DOSES OF MEDICINE.
10. Inspect family pets for signs of ringworm that should be treated by a veterinarian.



Administration	758-3820
Communicable Disease	758-3885
Community Resource Nurse	758-3675
Dental	758-3835
Environmental Health	758-3830
Family Planning	758-3870
Hearing Impaired Line	758-3725
Hearing & Vision	758-3890
HIV Testing & Counseling	758-3880
Immunization	758-3840
Laboratory	758-3825
Lead Testing	758-3768
Nursing Services	758-3850
Saginaw Health Plan	758-3855
Sexually Transmitted Diseases	758-3880
Substance Abuse Assessment	758-3781
After Hours	888-466-3141
WIC	758-3733



RINGWORM



Saginaw County
Department of Public Health

1600 N. Michigan Ave.
Saginaw, MI 48602
(989) 758-3800


“Protecting and promoting the public’s health
and well-being”
www.saginawpublichealth.org

Ringworm

Ringworm is actually not a worm at all! It is caused by a fungus (similar to molds and mushrooms) which is so small it can only be seen through a microscope. Ringworm fungus infects skin, hair, or nails. It is most commonly seen in children ages 2-20.



- ◆ Ringworm is **CONTAGIOUS** as long as lesions are present and live fungus remains in the lesions or on contaminated items.




- ◆ Ringworm grows best in moist places.

<u>Incubation Period</u>	
Scalp	10-14 days
Body	4-10 days

Mode of Spread

- ◆ Direct contact with the ringworm fungus from other people (skin to skin), animals, or soil.
- ◆ Indirectly by using items that have been contaminated by an infected person such as:



Scalp-dandruff-like scaling with patches of hair loss. Hair becomes brittle and breaks off easily either right at the scalp or slightly above the skin surface. There may be fluid or pus filled sores.




Body-starts out as tiny reddish, itchy, scaly bumps that spread into a ring-shaped sore. As the ring enlarges, the outside edges remain red and raised while the skin in the center of the ring takes on a more normal appearance.



Treatment

Scalp - Treatment is by prescription only. Consult your family physician.



Treatment usually consists of an anti-fungal medicine by mouth along with topical ointment and/or special shampoo. It takes **at least** four weeks to complete treatment.

Body - The preferred method of treatment is an anti-fungal cream prescribed by the doctor. It is usually applied once or twice a day.

- Over the counter anti-fungal cream may be used to treat ringworm on the body if prescription treatment is not available.

Caution: Do not use over the counter treatment on children two years of age and under. Use only with supervision on children 2-12 years of age.

- Treatment takes 2-4 weeks.

Exclusion from School

Children are excluded for 24 hours after beginning effective treatment. While under treatment and until sores are completely healed child is excluded from gym, swimming, and other activities likely to cause exposure to others such as contact sports.

